arXiv:1109.6812v1 [math.AP] 30 Sep 2011

COMPARABILITY AND REGULARITY ESTIMATES FOR SYMMETRIC NONLOCAL DIRICHLET FORMS

BARTŁOMIEJ DYDA AND MORITZ KASSMANN

Abstract. The aim of this work is to study comparability of nonlocal Dirichlet forms. We provide sufficient conditions on the kernel for local and global comparability. As an application we prove a-priori estimates in Hölder spaces for solutions to integrodifferential equations. These solutions are defined with the help of symmetric nonlocal Dirichlet forms.

1. Introduction

If, for every $x \in \mathbb{R}^d$, $A(x)$ is a positive definite matrix which is uniformly bounded, then for every ball $B \subset \mathbb{R}^d$ and every function $u \in C_c^\infty(B)$

$$\int_B \langle \nabla u(x), A(x) \nabla u(x) \rangle \, dx \asymp \int_B |\nabla u(x)|^2 \, dx \,. \tag{1}$$

This property is crucial for many questions related to partial differential operators of second order in divergence form and to diffusion processes generated by local Dirichlet forms. The aim of the present work is to study similar properties for symmetric nonlocal Dirichlet forms.

Fix $\alpha_0 \in (0, 2)$. Let $\mathcal{K}$ denote a family of kernels $k_\alpha : \mathbb{R}^d \times \mathbb{R}^d \to [0, \infty)$ which depend on indices $\alpha \in (\alpha_0, 2)$. We consider the corresponding bilinear forms

$$\iint_{\mathbb{R}^d \mathbb{R}^d} \big(u(y) - u(x)\big)\big(v(y) - v(x)\big) k_\alpha(x, y) \, dy \, dx \,, \qquad u, v \in C_c^\infty(\mathbb{R}^d) \,.$$

We study the question, under which additional assumptions on the kernels $k_\alpha \in \mathcal{K}$ local comparability holds, i.e. for every kernel $k_\alpha \in \mathcal{K}$, small ball B and every function $u \in C_c^\infty(B)$

$$\iint_{BB} \big(u(y) - u(x)\big)^2 k_\alpha(x, y) \, dy \, dx \asymp (2 - \alpha) \iint_{BB} \frac{\big(u(y) - u(x)\big)^2}{|x - y|^{d+\alpha}} \, dy \, dx \,. \tag{A}$$

This relation means that the ratio of the two quantities is bounded from below and above by two uniform positive constants which do not depend on $k_\alpha \in \mathcal{K}$.

Date: September 30, 2011.

2010 *Mathematics Subject Classification.* 31B05, 35B45, 35B05, 35R05, 47G20, 60J75.

Key words and phrases. Dirichlet forms, Hölder estimates.

Both authors have been supported by the German Science Foundation DFG through SFB 701. The first author was additionally supported by MNiSW grant N N201 397137.

Note that this problem is interesting and unsettled even if all kernels k_α correspond to one fixed index $\alpha \in (0,2)$. The case $\alpha = 2$ corresponds to (1).

For $\alpha \in (0,2)$ set $\mathcal{A}_{d,-\alpha} = \frac{\alpha\Gamma((d+\alpha)/2)}{2^{1-\alpha}\pi^{d/2}\Gamma(1-\alpha/2)}$. Note that $\mathcal{A}_{d,-\alpha} \asymp \alpha(2-\alpha)$ for all $\alpha \in (0,2)$. Fix $\alpha_0 \in (0,2)$ and $c > 0$. A standard example where relation (A) holds true is given by the family $\mathcal{K} = \{k_\alpha | \alpha \in (\alpha_0, 2)\}$ where k_α is any kernel satisfying

$$c\mathcal{A}_{d,-\alpha}|x-y|^{-d-\alpha} \leq k_\alpha(x,y) \leq c^{-1}\mathcal{A}_{d,-\alpha}|x-y|^{-d-\alpha} \tag{2}$$

for almost every $x, y \in \mathbb{R}^d$.

In this work we give sufficient conditions which are more general than (2). It is easy to see that (2) is not necessary for (A). Define $\widetilde{\mathcal{K}} = \{\widetilde{k}_\alpha | \alpha \in (\alpha_0, 2)\}$ with $\widetilde{k}_\alpha(x,y) = k_\alpha(x,y)(\mathbb{1}_{\{|x|\leq 0.1|y|\}} + \mathbb{1}_{\{|y|\leq 0.1|x|\}})$ where k_α is any kernel satisfying (2), then the kernels $\widetilde{k}_\alpha$ do not satisfy (2) but (A) is still satisfied for all $\widetilde{k}_\alpha \in \widetilde{\mathcal{K}}$.

One application of our investigation are local Poincaré- and Sobolev-inequalities, see [12, 14]. Those inequalities together with a class of appropriate cutoff-functions lead to regularity estimates for symmetric nonlocal Dirichlet forms. We assume that, for some constant $c > 0$, and every $R, \rho \in (0,1)$ there is a nonnegative function $\tau \in C^\infty(\mathbb{R}^d)$ with $\operatorname{supp}(\tau) = \overline{B_{R+\rho}}$, $\tau(x) \equiv 1$ on B_R, and for every k

$$\sup_{x\in\mathbb{R}^d} \int\limits_{\mathbb{R}^d} \big(\tau(y) - \tau(x)\big)^2 k(x,y)\, dy \leq c\rho^{-\alpha}\,. \tag{B}$$

Note that for $\alpha = 2$ Assumption (B) asks for the existence of a cut-off function τ with $\sup\limits_{x\in\mathbb{R}^d} |\nabla\tau|^2(x) \leq c\rho^{-2}$. Such τ obviously exists.

We are able to establish conditions (A) and (B) under quite mild assumptions. Let us always assume $k(x,y) = k(y,x)$ which is not a restriction since our bilinear forms are symmetric. Without mentioning it we always assume that for almost every $x, y \in \mathbb{R}^d$

$$L(x-y) \leq k(x,y) \leq U(x-y)\,, \tag{K}$$

for some functions $L, U : \mathbb{R}^d \to [0,\infty)$ satisfying $L(x) = L(-x)$, $U(x) = U(-x)$ for almost every $x \in \mathbb{R}^d$, $L \not\equiv 0$ on a set of positive measure, and

$$\int_{\mathbb{R}^d} (|z|^2 \wedge 1) U(z)\, dz \leq C_0 < \infty. \tag{U0}$$

Our main assumptions are the following:

(U1) There exists $C_1 > 0$ such that for every $r \in (0,1)$

$$\int_{B(0,r)} |z|^2 U(z)\, dz \leq C_1 r^{2-\alpha}\,. \tag{3}$$

(L1) There exist $a > 1$ and C_2, C_3 such that every annulus $B_{a^{-n+1}} \setminus B_{a^{-n}}$ ($n = 0, 1, \ldots$) contains a ball B_n with radius $C_2 a^{-n}$, such that

$$L(z) \geq C_3(2-\alpha)|z|^{-d-\alpha}, \quad z \in B_n. \tag{4}$$

Then we can prove the following result:

Theorem 1. *Assume the function* $k : \mathbb{R}^d \times \mathbb{R}^d \to [0,\infty)$ *satisfies (U0), (U1) and (L1). Then conditions (A) and (B) are satisfied. If the constants* C_0, C_1, C_2, C_3 *appearing in (U0), (U1), (L1) are independent of* $\alpha \in (\alpha_0, 2)$ $(\alpha_0 > 0)$, *then so are the constants in (A) and (B).*

Let us explain applications of this result. By $H^{\alpha/2}(\mathbb{R}^d)$ we denote the usual Sobolev space of fractional order $\alpha/2 \in (0,1)$, see (8). If $\Omega \subset \mathbb{R}^d$ is open and bounded, then $H^{\alpha/2}(\Omega)$ is the space of measurable functions $f : \Omega \to \mathbb{R}$ which can be represented as restrictions of $H^{\alpha/2}(\mathbb{R}^d)$ to Ω. $H^{\alpha/2}_{\text{loc}}(\Omega)$ denotes the space of all measurable functions $f : \Omega \to \mathbb{R}$ such that $\phi f \in H^{\alpha/2}(\Omega)$ for every $\phi \in C_c^\infty(\Omega)$.

Conditions (A) and (B) allow us to apply several techniques which were developed for partial differential operators of second order or local Dirichlet forms respectively. The following weak Harnack inequality holds true for supersolutions, see [10, 9].

Theorem 2. *Assume (A) and (B) hold true. Let* $\alpha_0 \in (0,2)$. *There are positive reals* p_0, c *such that for every* $\alpha \in (\alpha_0, 2)$ *and* $u \in L^\infty(\mathbb{R}^d) \cap H^{\alpha/2}_{loc}(B_1)$ *with* $u \geq 0$ *in* B_1 *satisfying* $\mathcal{E}(u,\phi) \geq 0$ *for every nonnegative* $\phi \in C_c^\infty(B_1)$ *the following inequality holds:*

$$c \inf_{B_{1/4}} u \geq \Big(\fint_{B_{1/2}} u(x)^{p_0}\, dx \Big)^{1/p_0} - c \sup_{x \in B_{1/2}} \int_{\mathbb{R}^d \setminus B_1} u^-(z) k(x,z)\, dz \,.$$

The constants p_0, c *depend only on* d, α_0 *and on the constants arising in (A) and (B).*

Throughout this article the abbreviation 'sup' shall denote the essential supremum and 'inf' the essential infimum. It is possible to combine Theorem 2 and Theorem 1 in order to obtain regularity estimates. In order to focus on the main issues we formulate a simple assumption on k for large values of $|x-y|$. We assume that there is $\gamma \in (0,\alpha)$ such that

$$\limsup_{R \to \infty} R^\gamma \int_{|z|>R} U(z)\, dz \leq 1 \,. \tag{U2}$$

Using conditions (L1), (U0), (U1) and (U2) the following nonlocal version of De-Giorgi's regularity result can be established [9].

Theorem 3. *Assume (L1), (U0), (U1) and (U2) hold true. Then there exist* $c > 0$, $\beta \in (0,1)$ *such that for every* $x_0 \in \mathbb{R}^n$, $u \in L^\infty(\mathbb{R}^d) \cap H^{\alpha/2}_{loc}(B_1(x_0))$ *satisfying* $\mathcal{E}(u,\phi) = 0$ *for every* $\phi \in C_c^\infty(B_1(x_0))$ *the following Hölder estimates holds for almost every* $x, y \in B_{1/2}(x_0)$:

$$|u(x) - u(y)| \leq c \|u\|_\infty |x-y|^\beta \,. \tag{5}$$

If the constants appearing in (L1), (U0), (U1) and (U2) are independent of $\alpha \in (\alpha_0, 2)$, *where* $\alpha_0 > 0$, *then so are the constants* c *and* β.

Theorem 3 is proved in resp. follows from the works [11, 2, 5, 3] if one allows the constant c in (5) to depend on $\alpha \in (0,2)$ with $c(\alpha) \to +\infty$ for $\alpha \to 2$ and if one imposes a stronger isotropic condition of the form

$$\forall\, z \in B_1(0): \quad L(z) \geq C_3 |z|^{-d-\alpha}.$$

Techniques which are robust as $\alpha \to 2$ are developed for equations in non-divergence form in [4].

The paper is organized as follows. In Section 2 we introduce some notation used in sequel and prove Theorem 1, breaking its proof into three parts, namely Propositions 5, 8 and 11. We also provide an example of a kernel satisfying (A) and (B), but not (L1), see Example 12. In Section 3 we provide the main ideas of how to prove Theorem 3.

2. Properties of the bilinear form

In this section we prove Theorem 1. The proof consists of several propositions and lemmata. At the end of the section we construct an example of a kernel satisfying (A) and (B), but not (L1).

Let us fix $\alpha \in (0,2)$ and consider the following quadratic forms

$$\mathcal{E}^k_D(u,u) = \int_D\int_D (u(y)-u(x))^2 k(x,y)\,dx\,dy, \quad u \in L^2(D), \tag{6}$$

$$\mathcal{E}^\alpha_D(u,u) = \alpha(2-\alpha)\int_D\int_D (u(y)-u(x))^2 |x-y|^{-d-\alpha}\,dx\,dy, \quad u \in L^2(D), \tag{7}$$

where $D \subset \mathbb{R}^d$ is some open set. Furthermore, we define Sobolev norm

$$\|u\|_{H^{\alpha/2}(\mathbb{R}^d)} = \|u\|_{L^2(\mathbb{R}^d)} + \mathcal{E}^\alpha_D(u,u)^{1/2}. \tag{8}$$

By $B(x,r) = \{z \in \mathbb{R}^d : |x-z| < r\}$ we denote the Euclidean ball with center x and radius $0 < r \leq \infty$, and we use an abbreviation $B_r = B(0,r)$. By $S^{d-1} = \{x \in \mathbb{R}^d : |x| = 1\}$ we denote the unit sphere.

We define Fourier transform as an isometry of $L^2(\mathbb{R}^d)$ determined by

$$\hat{u}(\xi) = (2\pi)^{-d/2}\int_{\mathbb{R}^d} u(x)e^{-i\xi\cdot x}\,dx, \quad u \in L^1(\mathbb{R}^d)\cap L^2(\mathbb{R}^d).$$

The following lemma contains a useful equivalent formulation of condition (U1).

Lemma 4. *Condition (U1) is equivalent to the following one:*

(U1') *There exists $C_4 > 0$ such that for every $r \in (0,1)$*

$$\int_{\mathbb{R}^d} \left(r^2 \wedge |z|^2\right) U(z)\,dz \leq C_4 r^{2-\alpha}. \tag{9}$$

If the constants C_0 and C_1 are independent of $\alpha \in (\alpha_0, 2)$, where $\alpha_0 > 0$, then so is the constant C_4, and vice versa.

Proof. Implication (U1') $\implies$ (U1) is obvious, we may take $C_0 = C_1 := C_4$. We assume now (U1) and we fix $0 < r < 1$. We consider $n = 0, 1, 2, \dots$ such that $2^{n+1} r < 1$ (the set of such n's is empty if $r \geq \frac{1}{2}$). We have by (U1)

$$\int_{2^n r<|z|\leq 2^{n+1}r} U(z)\,dz \leq 2^{-2n} r^{-2} \int_{2^n r<|z|\leq 2^{n+1}r} |z|^2 U(z)\,dz$$
$$\leq 2^{-2n} r^{-2} C_1\, 2^{(n+1)(2-\alpha)} r^{2-\alpha} = 2^{-n\alpha} 2^{2-\alpha} C_1 r^{-\alpha}.$$

After summing over all such n we obtain

$$\int_{r<|z|\leq 1/2} U(z)\,dz \leq \frac{2^{2-\alpha} C_1}{1-2^{-\alpha}} r^{-\alpha}.$$

Finally

$$\int_{1/2<|z|} U(z)\,dz \leq 4 \int_{\mathbb{R}^d} (|z|^2 \wedge 1) U(z)\,dz \leq 4C_0 \leq 4C_0 r^{-\alpha}.$$

Combining the two inequalities above and (U1) we get (U1') with $C_4 = (\frac{2^{2-\alpha}}{1-2^{-\alpha}} + 1)C_1 + 4C_0$. ☐

In next two propositions we prove the easier part of Theorem 1.

Proposition 5. *Condition (U1) implies (B). If the constants C_0 and C_1 are independent of $\alpha \in (\alpha_0, 2)$, where $\alpha_0 > 0$, then so is the constant in (B).*

Proof. Let $\tau \in C^\infty(\mathbb{R}^d)$ be a function satisfying $\mathrm{supp}(\tau) = \overline{B_{R+\rho}}$, $\tau \equiv 1$ on B_R, $0 \leq \tau \leq 1$ on $\mathbb{R}^d$ and $|\tau(x) - \tau(y)| \leq 2\rho^{-1}|x-y|$ for all $x, y \in \mathbb{R}^d$. In particular, we have then $|\tau(x) - \tau(y)| \leq (2\rho^{-1}|x-y|) \wedge 1$. For every $x \in \mathbb{R}^d$ we estimate, using Lemma 4

$$\int_{\mathbb{R}^d} (\tau(x) - \tau(y))^2 k(x,y)\,dy \leq \int_{\mathbb{R}^d} \left((4\rho^{-2}|x-y|^2) \wedge 1\right) U(x-y)\,dy$$
$$= 4\rho^{-2} \int_{\mathbb{R}^d} (|z|^2 \wedge \frac{\rho^2}{4}) U(z)\,dz$$
$$\leq 2^\alpha C_4 \rho^{-\alpha}.$$ ☐

In the proof of the next proposition we will need the following fact [8]. Its elementary proof may be found in [6], however one has to go through it and see that the constants do not depend on α, provided one has the factor $\alpha(2-\alpha)$ in front of the Gagliardo norm (7), (8).

Fact 6. *Let $D \subset \mathbb{R}^d$ be a bounded Lipschitz domain, and let $0 < \alpha < 2$. Then there exists a constant $c = c(d, D)$ (independent of α) and an extension operator $E : H^{\alpha/2}(D) \to H^{\alpha/2}(\mathbb{R}^d)$ with norm $\|E\| \leq c$.*

Furthermore, we will need the following Poincaré inequality [14].

Fact 7. *Let $D \subset \mathbb{R}^d$ be a bounded Lipschitz domain, and let $0 < \alpha_0 < \alpha < 2$. Then there exists a constant $c = c(d, \alpha_0, D)$ such that*

$$\|u - \frac{1}{|D|}\int_D u\,dx\|^2_{L^2(D)} \le c\mathcal{E}^\alpha_D(u,u), \quad u \in H^{\alpha/2}(D). \tag{10}$$

Now we are ready to formulate and prove the following comparability result.

Proposition 8. *Assume (U0), (U1) and let $0 < \alpha_0 < \alpha < 2$. If $D \subset \mathbb{R}^d$ is a bounded Lipschitz domain, then there exists a constant $c = c(\alpha_0, d, C_1, C_0, D)$ such that*

$$\mathcal{E}^k_D(u,u) \le c\mathcal{E}^\alpha_D(u,u), \quad u \in H^{\alpha/2}(D). \tag{11}$$

The constant c may be chosen such that (11) holds for all balls $D = B_r$ of radius $r < 1$, and for all $\alpha \in (\alpha_0, 2)$.

Proof. By E we denote the extension operator from $H^{\alpha/2}(D)$ to $H^{\alpha/2}(\mathbb{R}^d)$, see Fact 6. By subtracting a constant, we may and do assume that $\int_D u\,dx = 0$. We have by Plancherel formula and Fubini theorem

$$\begin{aligned}
\mathcal{E}^k_D(u,u) &\le \int_D\int_{D-y} (u(y+z) - u(y))^2 U(z)\,dz\,dy \qquad (12)\\
&\le \int_D\int_{B(0,\operatorname{diam} D)} (Eu(y+z) - Eu(y))^2 U(z)\,dz\,dy\\
&\le \int_{B(0,\operatorname{diam} D)}\int_{\mathbb{R}^d} (Eu(y+z) - Eu(y))^2\,dy\,U(z)\,dz\\
&= \int_{\mathbb{R}^d}\left(\int_{B(0,\operatorname{diam} D)} |e^{i\xi\cdot z} - 1|^2 U(z)\,dz\right)|\widehat{Eu}(\xi)|^2\,d\xi\\
&= \int_{\mathbb{R}^d}\left(\int_{B(0,\operatorname{diam} D)} 4\sin^2\Big(\frac{\xi\cdot z}{2}\Big) U(z)\,dz\right)|\widehat{Eu}(\xi)|^2\,d\xi. \qquad (13)
\end{aligned}$$

For $|\xi| > 2$ we obtain, using (U1')

$$\int 4\sin^2\Big(\frac{\xi\cdot z}{2}\Big)U(z)\,dz \le |\xi|^2\int (|z|^2 \wedge 4|\xi|^{-2})U(z)\,dz \le 4C_4|\xi|^\alpha, \tag{14}$$

and for $|\xi| \le 2$

$$\int 4\sin^2\Big(\frac{\xi\cdot z}{2}\Big)U(z)\,dz \le 4\int \left(\Big|\frac{\xi\cdot z}{2}\Big|^2 \wedge 1\right) U(z)\,dz \le 4C_0.$$

Thus

$$\begin{aligned}
\mathcal{E}^k_D(u,u) &\le c'\int_{\mathbb{R}^d} (|\xi|^\alpha + 1)\,|\widehat{Eu}(\xi)|^2\,d\xi\\
&\le c'\|Eu\|^2_{H^{\alpha/2}(\mathbb{R}^d)} \le c\|u\|^2_{H^{\alpha/2}(D)}\\
&= c(\mathcal{E}^\alpha_D(u,u) + \|u\|^2_{L^2(D)}) \qquad (15)
\end{aligned}$$

with $c = c(d, C_4, C_0, D)$. Since $\int_D u\,dx = 0$, we have by Fact 7

$$\mathcal{E}^\alpha_D(u,u) \ge c(\alpha_0, d, D)\int_D u^2(x)\,dx.$$

and this together with (15) proves (11).

By scaling, the last assertion of the Theorem is satisfied with a constant $c = c(\alpha_0, d, C_4, C_0, B_1)$. $\square$

The proof of the remaining part of Theorem 1, i.e. the inequality '$\geq$' in (A) under the assumption (L1), is more difficult. We will need the following two technical lemmata.

Lemma 9. *Let $0 < \alpha_0 < \alpha < 2$. We let $\eta \in (0,1)$ and for a ball $B = B(x,r)$ we denote $B^* = B(x, r/\eta)$. Suppose that for some c_k, $r_0 > 0$ and all $0 < r < r_0$ we have*

$$\mathcal{E}^k_{B^*}(u,u) \geq c_k \mathcal{E}^\alpha_B(u,u),$$

for every function u and every ball B of radius ηr. Then there exists a constant $c = c(d, \alpha_0, \eta)$, such that for every ball B of radius $r < r_0$ and every function u

$$\mathcal{E}^k_B(u,u) \geq c c_k \mathcal{E}^\alpha_B(u,u).$$

Proof. Fix some $0 < r < r_0$ and a ball D of radius r. We take $\mathcal{B}$ to be a family of balls with the following properties.

(i) For some $c = c(d)$ and any $x, y \in D$, if $|x - y| < c \operatorname{dist}(x, D^c)$, then there exists $B \in \mathcal{B}$ such that $x, y \in B$.
(ii) For every $B \in \mathcal{B}$, $B^* \subset D$.
(iii) Family $\{B^*\}_{B \in \mathcal{B}}$ has the finite overlapping property, that is, each point of D belongs to at most $M = M(d)$ balls B^*, where $B \in \mathcal{B}$.

Such a family $\mathcal{B}$ may be constructed by considering Whitney decomposition of D into cubes and then covering each Whitney cube by an appropriate family of balls.

We have

$$\begin{aligned}
\mathcal{E}^k_D(u,u) &\geq \frac{1}{M^2} \sum_{B \in \mathcal{B}} \int_{B^*} \int_{B^*} (u(x) - u(y))^2 k(x-y)\, dy\, dx \\
&\geq \frac{c_k}{M^2}(2-\alpha) \sum_{B \in \mathcal{B}} \int_B \int_B (u(x) - u(y))^2 |x-y|^{-d-\alpha}\, dy\, dx \\
&\geq \frac{c_k}{M^2}(2-\alpha) \int_D \int_{|x-y| < c \operatorname{dist}(x, D^c)} (u(x) - u(y))^2 |x-y|^{-d-\alpha}\, dy\, dx. \qquad (16)
\end{aligned}$$

By [7, Proposition 5 and proof of Theorem 1], we may estimate

$$\begin{aligned}
&\int_D \int_{|x-y| < c \operatorname{dist}(x, D^c)} (u(x) - u(y))^2 |x-y|^{-d-\alpha}\, dy\, dx \\
&\qquad \geq c(\alpha, d) \int_D \int_D (u(x) - u(y))^2 |x-y|^{-d-\alpha}\, dy\, dx, \qquad (17)
\end{aligned}$$

with some constant $c(\alpha, d)$. We note that in [7, proof of Theorem 1] the constant depends on the domain in question, but in our case, by scaling, we can take the same constant independent of the choice of the ball D. One may also check that $c(\alpha, d)$ stays bounded when $\alpha \in (\alpha_0, 2)$. By (16) and (17) the lemma follows. $\square$

Lemma 10. *If $q \in L^1(\mathbb{R}^d)$ is a nonnegative function with* $\operatorname{supp} q \subset B_\rho$*, then for all $R > 0$ and functions u*

$$\mathcal{E}^{q*q}_{B_R}(u,u) \le 4\|q\|_{L^1}\mathcal{E}^q_{B_{R+\rho}}(u,u).$$

Proof. We have

$$\begin{aligned}
\mathcal{E}^{q*q}_{B_R}(u,u) &= \int_{B_R}\int_{B_R}\int_{B_{R+\rho}} (u(x)-u(y))^2 q(x-z)q(z-y)\,dz\,dy\,dx \\
&\le 2\int_{B_R}\int_{B_R}\int_{B_{R+\rho}} \Big((u(x)-u(z))^2 + (u(z)-u(y))^2\Big) \\
&\qquad\qquad\qquad\qquad \times q(x-z)q(z-y)\,dz\,dy\,dx \\
&\le 4\int_{B_R}\int_{B_{R+\rho}} (u(x)-u(z))^2 q(x-z)\,dz\,dx \int q(y)\,dy \\
&\le 4\|q\|_{L^1}\mathcal{E}^q_{B_{R+\rho}}(u,u).
\end{aligned}$$

□

We are now ready to finish the proof of Theorem 1.

Proposition 11. *Assume that L satisfies (L1), and let $0 < \alpha_0 < \alpha < 2$. Then there exists a constant $c = c(d, \alpha_0, C_2, C_3, a)$, such that for all $0 < r < 1$*

$$\mathcal{E}^\alpha_{B_r}(u,u) \le c\mathcal{E}^L_{B_r}(u,u)$$

Proof. Let

$$q_n(z) = (L(z) \wedge C_3(2-\alpha)|z|^{-d-\alpha})\mathbb{1}_{B_{a^{-n}}\setminus B_{a^{-n-1}}}.$$

Using estimate $(a^\alpha - 1)/\alpha \le (a^2-1)/2$ it is easy to see that

$$\|q_n\|_{L^1} \le c(d, C_3, a)(2-\alpha)a^{n\alpha}.$$

Let $B_n \subset B_{a^{-n}} \setminus B_{a^{-n-1}}$ be a ball like in the assumption (L1), that is, having radius $C_2 a^{-n}$ and such that

$$L(z) \ge C_3(2-\alpha)|z|^{-d-\alpha} \ge C_3(2-\alpha)a^{(n+1)(d+\alpha)}, \quad z \in B_n \cup -B_n.$$

We obtain

$$\begin{aligned}
q_n * q_n(z) &\ge C_3^2(2-\alpha)^2 a^{2(n+1)(d+\alpha)}\,\mathbb{1}_{B_n\cup -B_n} * \mathbb{1}_{B_n \cup -B_n}(z) \\
&\ge (2-\alpha)^2 c(d, C_2, C_3, a)a^{nd+2n\alpha}\,\mathbb{1}_{B_{C_2a^{-n}}}(z).
\end{aligned}$$



FIGURE 1. Support of the kernel k from Example 12 (with $b = 1/2$) consists of four thorns. Also sets P_0 and P_1 (see (20)) are shown: four larger squares constitute set P_0, and four smaller – set P_1.

We fix $0 < r < 1$. Let n_0 be the smallest natural number such that $a^{-n_0} < r/2$. From inequality $L \geq \sum_{n=n_0}^{\infty} q_n$ and Lemma 10 we obtain

$$\begin{aligned}
\mathcal{E}^{L}_{B_r}(u,u) &\geq \sum_{n=n_0}^{\infty} \mathcal{E}^{q_n}_{B_{r/2+a^{-n}}}(u,u) \geq \sum_{n=n_0}^{\infty} (4\|q_n\|_{L^1})^{-1}\mathcal{E}^{q_n * q_n}_{B_{r/2}}(u,u) \\
&\geq c(d, C_2, C_3, a)(2-\alpha) \\
&\quad \times \sum_{n=n_0}^{\infty} \int_{B_{r/2}} \int_{B_{r/2}} (u(x)-u(y))^2 a^{n(d+\alpha)} \mathbb{1}_{B_{C_2 a^{-n}}}(x-y)\, dy\, dx \\
&\geq c'(d, C_2, C_3, a)(2-\alpha) \int_{B_{C_2 r/(4a)}} \int_{B_{C_2 r/(4a)}} \frac{(u(x)-u(y))^2}{|x-y|^{d+\alpha}}\, dy\, dx \\
&= c'(d, \alpha_0, C_2, C_3, a)\mathcal{E}^{\alpha}_{B_{C_2 r/(4a)}}(u,u).
\end{aligned}$$

The proof is finished by applying Lemma 9. □

Let us show that (L1) is not necessary for (A) and (B) to hold. The reason is that (A) uses only integrated quantities but not pointwise estimates on k. However, Assumption (L1) is weak and useful at the same time.

Example 12. Let $b \in (0,1)$ and

$$\Gamma = \{(x_1,x_2) \in \mathbb{R}^2 : |x_2| \geq |x_1|^b \text{ or } |x_1| \geq |x_2|^b\}.$$

We consider the following function

$$k(z) = (2-\alpha)\mathbb{1}_{\Gamma \cap B_1}(z)|z|^{-2-\beta}, \quad z \in \mathbb{R}^2, \tag{18}$$

where $\beta = \alpha - 1 + 1/b$, see Figure 1. We will show that for such a function k conditions (A) and (B) are satisfied.

We have, for $0 < r < 1$

$$\begin{aligned}\int_{B_r} |z|^2 k(z)\,dz &\leq 8(2-\alpha)\int_0^r \int_0^{x^{1/b}} (x^2+y^2)^{-\beta/2}\,dy\,dx \\ &\leq 8(2-\alpha)\int_0^r \int_0^{x^{1/b}} x^{-\beta}\,dy\,dx = 8r^{2-\alpha},\end{aligned} \tag{19}$$

hence k satisfies (U1) with $C_1 = 8$. For $n = 0,1,2,\ldots$, we consider the set

$$\begin{aligned}E_n =& \{(x_1,x_2) : 2^{-n-2} \leq |x_1| \leq 2^{-n-1} \text{ and } |x_2| \leq 2^{-2-(n+2)/b}\} \\ & \cup \{(x_1,x_2) : 2^{-n-2} \leq |x_2| \leq 2^{-n-1} \text{ and } |x_1| \leq 2^{-2-(n+2)/b}\}.\end{aligned}$$

We have $E_n \subset \Gamma$. Let

$$q_n(z) = 2^{n(\beta+2)}\mathbb{1}_{E_n}(z)$$

and

$$P_n = \{(x_1,x_2) : \frac{5}{4}2^{-n-2} \leq |x_1|, |x_2| \leq \frac{7}{4}2^{-n-2}\}. \tag{20}$$

If $(x_1,x_2) \in P_n$, then

$$\begin{aligned}q_n * q_n(x_1,x_2) &\geq \int_{-2^{-2-(n+2)/b}}^{2^{-2-(n+2)/b}} \int_{\{|z_2-x_2| \leq 2^{-2-(n+2)/b}\}} q_n(x_1-z_1, x_2-z_2)q_n(z_1,z_2)\,dz_2\,dz_1 \\ &= 2^{2n(\beta+2)}(2^{-1-(n+2)/b})^2 = 2^{-2-4/b}2^{n(2\alpha+2)}.\end{aligned} \tag{21}$$

We fix $R \in (0,1)$ and take the smallest natural number n_0 for which $2^{-n_0} < R/2$. Since $\|q_n\|_{L^1} = 2^{n\alpha-1-2/b} \leq 2^{n\alpha-2}$, from Lemma 10 we obtain

$$(2-\alpha)\sum_{n=n_0}^{\infty} 2^{-n\alpha}\mathcal{E}_{B_{R/2}}^{q_n * q_n}(u,u) \leq (2-\alpha)\sum_{n=n_0}^{\infty} \mathcal{E}_{B_R}^{q_n}(u,u) \leq \mathcal{E}_{B_R}^{k}(u,u). \tag{22}$$

On the other hand, by (21)

$$(2-\alpha)\sum_{n=n_0}^{\infty} 2^{-n\alpha} q_n * q_n(z) \geq (2-\alpha)2^{-2-4/b}\sum_{n=n_0}^{\infty} 2^{n(\alpha+2)}\mathbb{1}_{P_n}(z) =: f(z).$$

We note that each set $P_n \cap B_{2^{-n}} \setminus B_{2^{-n-1}}$ contains a ball B_{n+1} of radius $c2^{-n-1}$, where c is some universal constant. Furthermore, on this ball B_{n+1} we have

$$f(z) = (2-\alpha)2^{-2-4/b}2^{n(\alpha+2)} \geq (2-\alpha)2^{\alpha-4/b}|z|^{-(\alpha+2)} \geq (2-\alpha)2^{-4/b}|z|^{-(\alpha+2)},$$

provided $n \geq n_0$. Consequently, f satisfies (L1) with $a = 2$ and $n \geq n_0 + 1$, or, equivalently, a rescaled function $x \mapsto f(2^{-n_0-1}x)$ satisfies (L1) with $a = 2$. Since $R/8 < 2^{-n_0-1}$, we deduce from Proposition 11

$$\mathcal{E}^{\alpha}_{B_{R/8}}(u,u) \leq c\mathcal{E}^{f}_{B_{R/8}}(u,u) \leq 2^{2+4/b} c \sum_{n=n_0}^{\infty} 2^{-n\alpha} \mathcal{E}^{q_n * q_n}_{B_{R/2}}(u,u).$$

From this, (22) and Lemma 9 we deduce that

$$\mathcal{E}^{\alpha}_{B_R}(u,u) \leq c\mathcal{E}^{k}_{B_R}(u,u).$$

The reverse inequality follows from Proposition 8, hence (A) is satisfied. Also (B) is satisfied by Proposition 5.

3. Regularity estimates

In this section we provide the proof of Theorem 3. The main idea of the proof is to extend a result of [10].

Lemma 13. *Assume $x_0 \in \mathbb{R}^d$. For $r \in (0,1)$ and $x \in B_{r/2}(x_0)$ let ν^x_r be a measure on $\mathbb{R}^d \setminus B_r(x_0)$ satisfying*

$$\limsup_{j \to \infty} (\eta_{r,j})^{1/j} < 1, \qquad \text{where } \eta_{r,j} := \sup_{x \in B_{r/2}(x_0)} \nu^x_r(\mathbb{R}^d \setminus B_{2^j r}(x_0)) < \infty\,. \tag{23}$$

Assume that for some $c_1 \geq 1$, $p > 0$, every $r \in (0,1)$ and every $u \in L^\infty(\mathbb{R}^d) \cap H^{\alpha/2}_{loc}(B_r(x_0))$ satisfying $\mathcal{E}(u,\phi) = 0$ for every $\phi \in C^\infty_c(B_r(x_0))$ and $u \geq 0$ in $B_r(x_0)$, the weak Harnack inequality

$$\Big(\fint_{B_{r/2}(x_0)} u(x)^p \, dx \Big)^{1/p} \leq c_1 \inf_{x \in B_{r/4}(x_0)} u + c_1 \sup_{x \in B_{r/2}(x_0)} \int_{\mathbb{R}^d} u^-(z) \nu^x_r(dz)\,. \tag{24}$$

holds true. Then there exist $\beta \in (0,1)$, $c > 0$ such that for every $r \in (0,1)$ and every $u \in L^\infty(\mathbb{R}^d) \cap H^{\alpha/2}_{loc}(B_r(x_0))$ satisfying $\mathcal{E}(u,\phi) = 0$ for every $\phi \in C^\infty_c(B_r(x_0))$ and every $\rho \in (0, r/2)$ the following regularity estimate holds:

$$\sup_{x,y \in B_\rho(x_0)} |u(x) - u(y)| \leq c\|u\|_\infty (\rho/r)^\beta\,. \tag{25}$$

If c_1, p and the limes superior in (23) are independent of x_0 or α, then so is c.

Remark 1. Instead of (24) one may assume the strong Harnack inequality

$$\sup_{x \in B_{r/4}(x_0)} u \leq c_1 \inf_{x \in B_{r/4}(x_0)} u + c_1 \sup_{x \in B_{r/2}(x_0)} \int_{\mathbb{R}^d} u^-(z) \nu^x_r(dz)\,. \tag{26}$$

One only needs to change the constant c_2 in the proof of the lemma.

Proof. The idea is to adopt the methods of [13] to the nonlocal situation, see also [15]. Fix $x_0 \in \mathbb{R}^d$. Let $c_1 > 0$ be the constant in (24). Let $\theta = 4$. Set $c_2 = c_1 \theta^{d/p} 2^{\frac{1-d}{p}}$ and $\kappa = (c_2)^{-1}/2$. Let $\beta \in (0, \ln(\frac{2}{2-\kappa})/\ln(\theta))$ be another constant to be fixed later. Note that $(1 - \frac{\kappa}{2}) \leq \theta^{-\beta}$.

Let $r > 0$ and $u \in L^\infty(\mathbb{R}^d) \cap H^{\alpha/2}_{\text{loc}}(B_r(x_0))$ satisfy $\mathcal{E}(u,\phi) = 0$ for every $\phi \in C_c^\infty(B_r(x_0))$. We can assume $u(x_0) = 0$ which can be obtained by adding a constant if needed. Let us write B_r instead of $B_r(x_0)$ for $r > 0$.

We will construct an increasing sequence (m_n) and a decreasing sequence (M_n) satisfying for every $n \in \mathbb{Z}$

$$\begin{aligned} &m_n \le u(x) \le M_n \quad \text{for almost all } x \in B_{r\theta^{-n}}\,, \\ &M_n - m_n \le K\theta^{-n\beta}\,, \end{aligned} \tag{27}$$

where $K = M_0 - m_0 \in [0, 2\|u\|_\infty]$. Set $M_0 = \|u\|_\infty$, $m_0 = \inf\limits_{\mathbb{R}^d} u(x)$ and $M_{-n} = M_0$, $m_{-n} = m_0$ for every $n \in \mathbb{N}$. Assume there is $k \in \mathbb{N}$ and there are M_n, m_n such that (27) holds for $n \le k-1$. We need to choose m_k, M_k such that (27) holds for $n = k$.

For $x \in \mathbb{R}^d$ set

$$v(x) = \Big(u(x) - \frac{M_{k-1} + m_{k-1}}{2}\Big)\frac{2\theta^{(k-1)\beta}}{K}\,,$$

The definition of v implies $|v(x)| \le 1$ for almost every $x \in B_{r\theta^{-(k-1)}}$ and $\mathcal{E}(v,\phi) = 0$ for every $\phi \in C_c^\infty(B_r)$.

We now derive a pointwise estimate of the function v on $\mathbb{R}^d \setminus B_{r\theta^{-(k-1)}}$. Given $y \in \mathbb{R}^d$ with $|y - x_0| \ge r\theta^{-(k-1)}$ there is $j \in \mathbb{N}$ such that

$$r\theta^{-k+j} \le |y - x_0| < r\theta^{-k+j+1}\,.$$

For such y and j we conclude

$$\begin{aligned} \frac{K}{2\theta^{(k-1)\beta}} v(y) &= \Big(u(y) - \frac{M_{k-1} + m_{k-1}}{2}\Big) \\ &\le \Big(M_{k-j-1} - m_{k-j-1} + m_{k-j-1} - \frac{M_{k-1} + m_{k-1}}{2}\Big) \\ &\le \Big(M_{k-j-1} - m_{k-j-1} - \frac{M_{k-1} - m_{k-1}}{2}\Big) \\ &\le \Big(K\theta^{-(k-j-1)\beta} - \tfrac{K}{2}\theta^{-(k-1)\beta}\Big)\,, \\ \text{i.e. } v(y) &\le 2\theta^{j\beta} - 1 \ \le 2\Big(\theta\frac{|y-z|}{r\theta^{-(k-1)}}\Big)^\beta - 1\,. \end{aligned}$$

Analogously,

$$v(y) \ge 1 - 2\theta^{j\beta} \ \ge 1 - 2\Big(\theta\frac{|y-z|}{r\theta^{-(k-1)}}\Big)^\beta\,.$$

Now there are two cases:

Case 1: $|\{x \in B_{r\theta^{-k}} : v(x) \le 0\}| \ge \frac{1}{2}|B_{r\theta^{-k}}|$

Case 2: $|\{x \in B_{r\theta^{-k}} : v(x) > 0\}) \ge \frac{1}{2}|B_{r\theta^{-k}}|$

We work out details for Case 1 and comment afterwards on Case 2. In Case 1 our aim is to show $v(x) \le 1 - \kappa$ for almost every $x \in B_{r\theta^{-k}}$. Because then for almost every $x \in B_{r\theta^{-k}}$

$$\begin{aligned} u(x) &\leq \frac{M_{k-1} + m_{k-1}}{2} + \tfrac{(1-\kappa)K}{2}\theta^{-(k-1)\beta} \\ &= m_{k-1} + \frac{M_{k-1} - m_{k-1}}{2} + \tfrac{(1-\kappa)K}{2}\theta^{-(k-1)\beta} \\ &\leq m_{k-1} + \tfrac{K}{2}\theta^{-(k-1)\beta} + \tfrac{(1-\kappa)K}{2}\theta^{-(k-1)\beta} = m_{k-1} + (1 - \tfrac{\kappa}{2})K\theta^{-(k-1)\beta} \\ &\leq m_{k-1} + K\theta^{-k\beta}\,. \end{aligned} \tag{28}$$

In this case we set $m_k = m_{k-1}$ and $M_k = m_k + K\theta^{-k\beta}$ and obtain, using (28), $m_k \leq u(x) \leq M_k$ for almost every $x \in B_{r\theta^{-k}}$, what needs to be proved.

Let us show $v(x) \leq 1 - \kappa$ for almost every $x \in B_{r\theta^{-k}}$. Consider $w = 1 - v$. Then $\mathcal{E}(w, \phi) = 0$ for every $\phi \in C_c^\infty(B_{r\theta^{-k+1}})$ and $w \geq 0$ in $B_{r\theta^{-k+1}}$. We apply (24) and obtain

$$\Big(\fint\limits_{B_{\frac{1}{2}r\theta^{-k+1}}} w(x)^p \, dx \Big)^{1/p} \leq c_1 \inf_{B_{\frac{1}{4}r\theta^{-k+1}}} w + c_1 \sup_{x \in B_{\frac{1}{2}r\theta^{-k+1}}} \int\limits_{\mathbb{R}^d} w^-(z) \nu^x_{r\theta^{-k+1}}(dz)\,. \tag{29}$$

In the situation of Case 1 we obtain

$$(1/2)^{1/p} \leq \Big(\fint\limits_{B_{r\theta^{-k}}} w(x)^p \, dx \Big)^{1/p} \leq (\tfrac{\theta}{2})^{d/p} \Big(\fint\limits_{B_{\frac{1}{2}r\theta^{-k+1}}} w(x)^p \, dx \Big)^{1/p} \tag{30}$$

$$\leq c_1 (\tfrac{\theta}{2})^{d/p} \inf_{B_{r\theta^{-k}}} w + c_1 (\tfrac{\theta}{2})^{d/p} \sup_{x \in B_{\frac{1}{2}r\theta^{-k+1}}} \int\limits_{\mathbb{R}^d} w^-(y) \nu^x_{r\theta^{-(k-1)}}(dy)\,. \tag{31}$$

For $0 < R < S$ let us abbreviate the annulus $B_S(x_0) \setminus B_R(x_0)$ by $A_{R,S}(x_0)$. Then we obtain

$$\begin{aligned} \inf_{B_{r\theta^{-k}}} w &\geq (c_2)^{-1} - \sup_{x \in B_{\frac{1}{2}r\theta^{-k+1}}} \int\limits_{\mathbb{R}^d} w^-(y) \nu^x_{r\theta^{-(k-1)}}(dy) \\ &\geq (c_2)^{-1} - \sum_{j=1}^\infty \sup_{x \in B_{\frac{1}{2}r\theta^{-k+1}}} \int\limits_{\mathbb{R}^d} \mathbb{1}_{A_{r\theta^{-k+j}, r\theta^{-k+j+1}}(x_0)} (1 - v(y))^- \, \nu^x_{r\theta^{-(k-1)}}(dy)\,, \\ &\geq (c_2)^{-1} - \sum_{j=1}^\infty (2\theta^{j\beta} - 2) \eta_{r\theta^{-(k-1)}, 2(j-1)} \\ &= (c_2)^{-1} - 2\sum_{j=1}^\infty (\theta^{j\beta} - 1) \eta_{r\theta^{-(k-1)}, 2(j-1)}\,. \end{aligned}$$

Assumption (23) guarantees $\sum\limits_{j=1}^\infty \theta^{j\beta} \eta_{r\theta^{-(k-1)}, 2(j-1)} < \infty$ if $0 < \beta < \beta_0$ and β_0 is sufficiently small. Choose β_0 accordingly. Then there is $l \in \mathbb{N}$ with

$$\sum_{j=l+1}^\infty (\theta^{j\beta_0} - 1) \eta_{r\theta^{-(k-1)}, 2(j-1)} \leq \sum_{j=l+1}^\infty \theta^{j\beta_0} \eta_{r\theta^{-(k-1)}, 2(j-1)} \leq (8c_2)^{-1}\,.$$

Given l we choose $\beta \in (0, \beta_0)$ sufficiently small such that

$$\sum_{j=1}^{l} (\theta^{j\beta} - 1)\eta_{r\theta^{-(k-1)},2(j-1)} \leq (8c_2)^{-1}\,.$$

Thus $w \geq \kappa$ on $B_{r\theta^{-k}}$ or , equivalently, $v \leq 1 - \kappa$ on $B_{r\theta^{-k}}$.

In Case 2 our aim is to show $v(x) \geq -1 + \kappa$. This time, set $w = 1 + v$. Following the strategy above one sets $M_k = M_{k-1}$ and $m_k = M_k - K\theta^{-k\beta}$ leading to the desired result.

Let us show how (27) proves the assertion of the lemma. Let $\rho \in (0, r/2)$. Choose $m \in \mathbb{N}_0$ with $r\theta^{-(m+1)} \leq \rho < r\theta^{-m}$. Then condition (27) implies

$$\sup_{x,y \in B_\rho(x_0)} |u(x) - u(y)| \leq K\theta^{-m\beta} = (r\theta^{-m-1})^\beta r^{-\beta} K\theta^\beta \leq K\theta^\beta \Big(\frac{\rho}{r}\Big)^\beta\,.$$

The assertion of the lemma follows and the proof is complete. □

Let us explain the proof of our main application.

Proof of Theorem 3. The proof of Theorem 3 follows from Lemma 13 if we can show that, for every $r \in (0,1)$ and every $u \in L^\infty(\mathbb{R}^d) \cap H^{\alpha/2}_{\mathrm{loc}}(B_r(x_0))$ satisfying $\mathcal{E}(u, \phi) = 0$ for every $\phi \in C_c^\infty(B_r(x_0))$ and $u \geq 0$ in $B_r(x_0)$, the weak Harnack inequality (24) holds true with $(\nu_r^x)_{x \in B_{r/2}(x_0)}$ satisfying (23).

Fix $x_0 \in \mathbb{R}^d$. Note that none of the constants below will depend on x_0. For $r \in (0,1)$ and $x \in B_{r/2}(x_0)$ define a measure ν_r^x on $\mathbb{R}^d \setminus B_r(x_0)$ by

$$\nu_r^x(A) = \int_A U(y - x)\, dy \,\Big(\int_{\mathbb{R}^d \setminus B_r(x_0)} U(y - x_0)\, dy \Big)^{-1}$$

for every Borel set $A \subset \mathbb{R}^d \setminus B_r(x_0)$. Assumption (U2) implies that there are $c_1 > 0$ and $R_0 > 1$ such that for every $R > R_0$, $r \in (0,1)$ and $x \in B_{r/2}(x_0)$

$$\int_{\mathbb{R}^d \setminus B_R(x_0)} U(z - x)\, dz \leq c_1 R^{-\gamma} \tag{32}$$

Because of Assumptions (K) and (L1) there is $c_2 > 0$ with

$$\Big(\int_{\mathbb{R}^d \setminus B_r} U(z)\, dz \Big)^{-1} \leq \Big(\int_{\mathbb{R}^d \setminus B_r} L(z)\, dz \Big)^{-1} \leq c_2 C_3 r^\alpha\,. \tag{33}$$

Estimates (32) and (33) imply:

$$\exists c_3 \geq 1\ \forall r \in (0,1)\ \exists j_0 \geq 1\ \forall j \geq j_0\ \forall x \in B_{r/2}(x_0):$$
$$\nu_r^x\big(\mathbb{R}^d \setminus B_{2^j r}(x_0)\big) \leq c_3 (2^j r)^{-\gamma} / r^{-\alpha} \leq c_3 2^{-\gamma j}\,.$$

Recall that we assume $\gamma < \alpha$ in (U2). Condition (23) now holds true because of $2^{-\gamma} < 1$ and $c_3^{1/j} \to 1$ for $j \to \infty$.

Let $r \in (0,1)$ and $u \in L^\infty(\mathbb{R}^d) \cap H^{\alpha/2}_{\mathrm{loc}}(B_r(x_0))$ satisfy $\mathcal{E}(u,\phi) = 0$ for every $\phi \in C_c^\infty(B_r(x_0))$ and $u \geq 0$ in $B_r(x_0)$. Then Theorem 2 implies

$$c_4 \inf_{B_{r/4}(x_0)} u \geq \Big(\fint_{B_{r/2}(x_0)} u(x)^{p_0}\, dx \Big)^{1/p_0} - c_4 \sup_{x\in B_{r/2}(x_0)} r^\alpha \int_{\mathbb{R}^d \setminus B_r(x_0)} u^-(z) k(x,z)\, dz$$

with some appropriate constant $c_4 > 0$. Here we replaced the radius One by some arbitrary radii $r \in (0,1)$. This is possible since $(L1)$, $(U1)$ and $(U2)$ allow for scaling. Finally, we note that, with some $c_5 > 0$

$$\begin{aligned}
&\sup_{x\in B_{r/2}(x_0)} r^\alpha \int_{\mathbb{R}^d\setminus B_r(x_0)} u^-(z) k(x,z)\, dz \\
&\leq c_5 \Big(\int_{\mathbb{R}^d\setminus B_r(x_0)} U(y - x_0)\, dy \Big)^{-1} \sup_{x\in B_{r/2}(x_0)} \int_{\mathbb{R}^d\setminus B_r(x_0)} u^-(z) U(z-x)\, dz\,,
\end{aligned}$$

where we used Assumption (K) and the estimate

$$\int_{\mathbb{R}^d\setminus B_r} U(z)\, dz \leq r^{-2} \int_{\mathbb{R}^d\setminus B_r} (r^2 \wedge |z|^2) U(z)\, dz \leq C_4 r^{-\alpha}\,, \tag{34}$$

which follows from Lemma 4. Condition (24) now follows. The proof is complete. □

4. Appendix

In this appendix we provide a global comparability result, i.e. we study comparability in the whole of $\mathbb{R}^d$.

Proposition 14. *If (U0) and (U1) hold, then there exists a constant* $c = c(\alpha, d, C_1, C_0)$ *such that*

$$\mathcal{E}^k_{\mathbb{R}^d}(u,u) \leq c(\mathcal{E}^\alpha_{\mathbb{R}^d}(u,u) + \|u\|^2_{L^2(\mathbb{R}^d)}), \quad u \in L^2(\mathbb{R}^d). \tag{35}$$

Furthermore, if (3) is satisfied for all $r > 0$*, then*

$$\mathcal{E}^k_{\mathbb{R}^d}(u,u) \leq c\mathcal{E}^\alpha_{\mathbb{R}^d}(u,u), \quad u \in L^2(\mathbb{R}^d). \tag{36}$$

If the constants C_0 and C_1 in (U0) and (U1) are independent of $\alpha \in (\alpha_0, 2)$, where $\alpha_0 > 0$, then so are the constants in (35) *and* (36)*.*

Proof. By E we denote the identity operator from $H^{\alpha/2}(\mathbb{R}^d)$ to itself. One easily checks that the proof of Proposition 8 from (12) until (15) works also in the present case of $D = \mathbb{R}^d$. Hence (35) follows.

To prove (36) we first observe that when (3) holds for all $r > 0$, then we may also get inequality (9) in Lemma 4 for *all* $r > 0$. Consequently, (14) holds for *all* $r > 0$, we plug it into (13) and we are done. □

We consider the following condition.

(K2,r_0) There exists $c_0 > 0$ such that for all $h \in S^{d-1}$ and all $0 < r < r_0$

$$\int_{\mathbb{R}^d} r^2 \sin^2\Big(\frac{h\cdot z}{r}\Big) L(z)\,dz \geq c_0 r^{2-\alpha}. \tag{37}$$

Clearly (L1) implies (K2,r_0) for $r_0 = 1$, and if C_1 is independent of $\alpha \in (\alpha_0, 2)$, where $\alpha_0 > 0$, then so is c_0. Condition (K2,r_0) is also satisfied if for all $h \in S^{d-1}$ and all $0 < r < r_0$

$$\int_{B(0,r)} |h\cdot z|^2 L(z)\,dz \geq c_2 r^{2-\alpha}. \tag{38}$$

We note that (39) under condition (38) has been proved in [1] by Abels and Husseini. The following theorem extends their result by giving a *characterisation* of functions L admitting comparability (39). We stress that $r_0 = \infty$ is allowed, and in such a case we put $\frac{1}{r_0^\alpha} = 0$.

Theorem 15. *Let* $0 < r_0 \leq \infty$*. If (K2,*r_0*) holds, then*

$$\mathcal{E}^\alpha_{\mathbb{R}^d}(u,u) \leq \frac{1}{c_0}\mathcal{E}^k_{\mathbb{R}^d}(u,u) + \frac{2^\alpha}{r_0^\alpha}\|u\|^2_{L^2}, \quad u \in C^1_c(\mathbb{R}^d). \tag{39}$$

Conversely, if for some $c < \infty$

$$\mathcal{E}^\alpha_{\mathbb{R}^d}(u,u) \leq c \iint (u(x)-u(y))^2 L(x-y)\,dy\,dx + \frac{2^\alpha}{r_0^\alpha}\|u\|^2_{L^2}, \quad u \in \mathcal{S}(\mathbb{R}^d), \tag{40}$$

then (K2,r_0*) holds.*

Proof. We change the variable x to $y+z$ and use Plancherel formula. Recalling that $(u(\cdot+z))^\wedge(\xi) = e^{i\xi\cdot z}\hat{u}(\xi)$ we obtain

$$\begin{aligned}
\mathcal{E}^k_{\mathbb{R}^d}(u,u) &\geq \iint (u(x)-u(y))^2 L(x-y)\,dy\,dx \\
&= \int \Big(\int |e^{i\xi\cdot z} - 1|^2 L(z)\,dz\Big) |\hat{u}(\xi)|^2\,d\xi \\
&= \int \Big(\int 4\sin^2\Big(\frac{\xi\cdot z}{2}\Big) L(z)\,dz\Big) |\hat{u}(\xi)|^2\,d\xi. \qquad (41)
\end{aligned}$$

If (K2,r_0) holds, then for all $|\xi| > 2/r_0$

$$\int 4\sin^2\Big(\frac{\xi\cdot z}{2}\Big) L(z)\,dz \geq \frac{4c_0}{2^\alpha}|\xi|^\alpha \geq c_0|\xi|^\alpha.$$

For $|\xi| \leq 2/r_0$ we have $|\xi|^\alpha \leq (2/r_0)^\alpha$. Inequality (39) follows from

$$\frac{\mathcal{A}_{d,-\alpha}}{2\alpha(2-\alpha)}\mathcal{E}^\alpha_{\mathbb{R}^d}(u,u) = \int_{\mathbb{R}^d} |\xi|^\alpha |\hat{u}(\xi)|^2\,d\xi. \tag{42}$$

Now we prove the converse. Assume (40). By (41), the right hand side of (40) equals

$$\int \Big(c\int 4\sin^2\Big(\frac{\xi\cdot z}{2}\Big) L(z)\,dz + \frac{2^\alpha}{r_0^\alpha}\Big) |\hat{u}(\xi)|^2\,d\xi,$$

hence by (42) and (40) we obtain that

$$c\int 4\sin^2\Big(\frac{\xi\cdot z}{2}\Big)L(z)\,dz+\frac{2^\alpha}{r_0^\alpha}\geq|\xi|^\alpha,\quad \text{for a.e. } \xi\in\mathbb{R}^d. \tag{43}$$

By continuity of the function

$$\mathbb{R}^d\setminus\{0\}\ni\xi\mapsto\int 4\sin^2\Big(\frac{\xi\cdot z}{2}\Big)L(z)\,dz,$$

(43) holds for all $\xi\in\mathbb{R}^d$. For $|\xi|\geq 2^{1+1/\alpha}r_0^{-1}$ we have by (43)

$$c\int 4\sin^2\Big(\frac{\xi\cdot z}{2}\Big)L(z)\,dz\geq\frac{|\xi|^\alpha}{2},$$

and hence (K2,$2^{-1/\alpha}r_0$) holds with $c_0=2^{\alpha-3}c^{-1}$. Since

$$\sin^2\Big(\frac{h\cdot z}{2r}\Big)\geq\frac{1}{4}\sin^2\Big(\frac{h\cdot z}{r}\Big),$$

also (K2,r_0) holds with *some* constant c_0. □

References

[1] H. Abels and R. Husseini. On hypoellipticity of generators of Lévy processes.

[2] R. F. Bass and D. A. Levin. Transition probabilities for symmetric jump processes. *Trans. Amer. Math. Soc.*, 354(7):2933–2953 (electronic), 2002.

[3] L. Caffarelli, C.-H. Chan, and A. Vasseur. Regularity theory for parabolic nonlinear integral operators. Preprint.

[4] L. Caffarelli and L. Silvestre. Regularity theory for fully nonlinear integro-differential equations. *Comm. Pure Appl. Math.*, 62(5):597–638, 2009.

[5] Z.-Q. Chen and T. Kumagai. Heat kernel estimates for stable-like processes on d-sets. *Stochastic Process. Appl.*, 108(1):27–62, 2003.

[6] E. Di Nezza, G. Palatucci, and E. Valdinoci. Hitchhiker's guide to the fractional Sobolev spaces. arXiv:1104.4345v2 [math.FA], 2011.

[7] B. Dyda. On comparability of integral forms. *J. Math. Anal. Appl.*, 318(2):564–577, 2006.

[8] D. Jerison and C. E. Kenig. The inhomogeneous Dirichlet problem in Lipschitz domains. *J. Funct. Anal.*, 130(1):161–219, 1995.

[9] M. Kassmann. Harnack's inequality and Hoelder regularity estimates for nonlocal operators. Preprint.

[10] M. Kassmann. A priori estimates for integro-differential operators with measurable kernels. *Calc. Var. Partial Differential Equations*, 34(1):1–21, 2009.

[11] T. Komatsu. Uniform estimates for fundamental solutions associated with non-local Dirichlet forms. *Osaka J. Math.*, 32(4):833–860, 1995.

[12] V. Maz′ya and T. Shaposhnikova. On the Bourgain, Brezis, and Mironescu theorem concerning limiting embeddings of fractional Sobolev spaces. *J. Funct. Anal.*, 195(2):230–238, 2002.

[13] J. Moser. On Harnack's theorem for elliptic differential equations. *Comm. Pure Appl. Math.*, 14:577–591, 1961.

[14] A. C. Ponce. An estimate in the spirit of Poincaré's inequality. *J. Eur. Math. Soc. (JEMS)*, 6(1):1–15, 2004.

[15] L. Silvestre. Hölder estimates for solutions of integro-differential equations like the fractional Laplace. *Indiana Univ. Math. J.*, 55(3):1155–1174, 2006.

FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT BIELEFELD, POSTFACH 100131, D-33501 BIELEFELD